

Suicide in Missouri: Youth Ages 10-24


University of Missouri—St. Louis

July 2015 - Brief No. 5


Suicide is the second leading cause of death among 10-24 year old youth. In 2013, 118 youth died by suicide.

Males take their lives at a dramatically higher rate than females.

Almost half of all youth suicides in 2013 involved firearms.




- Suicide is the second leading cause of death among 10-24 year olds in Missouri, exceeded only by unintentional injury (including motor vehicle accidents). In 2013, 310 youth died by unintentional injury, and 118 died by suicide.
- Most (75%) of youth who died by suicide in 2013 were 18-24.
- Among all youth suicides in 2013, the highest percentage was by white males (68%).
- No African American female youth died by suicide.



- Youth suicide rates in Missouri have remained fairly stable over time and are higher than rates nationally (8.8 per 100,000 compared to 7.3 from 2003 to 2013).
- The largest *number* of youth suicides occurred in the St. Louis and Kansas City metropolitan areas. Counties with the highest suicide *rates* are in rural Missouri.

Means


Means of Suicide among Missouri Youth Ages 10-24 (2009-2013)




- From 2009-2013, an average of 50% of all youth suicides involved firearms.
- During that same time period, an average of 38% of youth used some means of suffocation.
- Among males 10-17, suffocation was the primary means of suicide, followed by firearms. Firearm use was the primary means of suicide among males ages 18-24.

- Older male youth (18-24) used firearms far more than females in the same age group to take their lives (53% compared to 38%).
- Females 10-17 used suffocation more often than other means. Among female youth 18-24, suffocation and firearms were used most frequently.
- Among females 18-24, 19% ended their lives by poisoning themselves, compared to only 5% of males. No youth under 18 killed themselves using poison.

Means of Suicide among Missouri Youth by Sex (2009-2013)




Rates of Youth Suicide by Means of Death: Missouri Youth Ages 10-24 (2009-2013)




- The means by which youth take their lives has remained relatively steady over time.
- Suicide rates are higher for those who use firearms compared to those who use suffocation.

Intentional Self-Injury*




- Among youth 10-19, intentional self-injury rates are higher among females than males. However, there are almost no gender differences for youth 20-24.
- While the *suicide* rate for young males is five times that for females, the *intentional self-injury* rate is higher among females.
- In 2012, youth aged 15-17 years old had the highest reported intentional self-injury rates of any age group followed by 18-19 year old youth.




*Intentional self-injury data are from the Department of Health and Senior Services (DHSS) Missouri Information for Community Assessment (MICA) and were obtained from in-patient hospital or emergency room records of individuals treated for intentional self-injuries.

Suicidal Ideation



- According to the 2014 Missouri Student Survey, 12% of middle and high school students reported seriously considering suicide in the year prior to the survey, 9% said they had made a suicide plan and 5% said they attempted suicide. Of those who attempted suicide, 3% were male and 7% were female.
- While more females than males seriously considered, planned or attempted suicide, attempts requiring medical attention were more similar between males and females.
- White students were less likely than students of any other racial/ethnic group to report that they had attempted suicide in the past year.
- Suicidal ideation and suicide attempts were highest among 8th grade students. The percentage reporting they needed a doctor or nurse’s care for a suicide attempt was similar among younger and older youth.



Suicide data are from the Department of Health and Senior Services (DHSS) Missouri Information for Community Assessment (MICA) with the following exceptions: 1) means data from 2009-2013 and Missouri/U.S. comparison data are from the Centers for Disease Control Web-based Injury Statistics Query and Reporting System (WISQARS); and 2) data on suicidal ideation are from the 2014 Missouri Student Survey. “Firearms” include pistols, rifles, and shotguns; “Suffocation” includes hanging and asphyxiation; “Poisoning” includes drug overdoses; and “Other” includes jumping, cutting, drowning, and unspecified/other.

Missouri Institute of Mental Health at the University of Missouri-St. Louis

This brief was produced by researchers at MIMH for the Missouri Suicide Prevention Project, a joint effort between the Missouri Department of Mental Health and MIMH. Production of this brief was made possible by grant number SM060377 from the Substance Abuse and Mental Health Services Administration (SAMHSA). The views, policies, and opinions expressed are those of the authors and do not necessarily reflect those of SAMHSA. For questions or more information about these data email Missouri Suicide Prevention Project at MSPP@dmh.mo.gov.